

# β-actin Monoclonal Antibody(5B7), Biotin Conjugated

CatalogNo: YM2192

## Key Features

### Host Species

- Mouse

### Reactivity

- Human,Rat,Mouse,Mk,Dg,Ch,Hamster,Rabbit,Insect

### Applications

- IF,WB,IHC

### Isotype

- IgG1

### Conjugate

- Biotin

## Recommended Dilution Ratios

Optimal working dilutions should be determined experimentally by the investigator

Suggested starting dilutions are as follows:WB 1:5000

IHC 1:200.

## Storage

### Storage\*

Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.

### Formulation

1mg/ml

## Basic Information

### Clonality

Monoclonal

### Clone Number

5B7

## Immunogen Information

### Specificity

β-actin Monoclonal Antibody(5B7) Biotin conjugated specially designed for your WB or IHC analysis.

## Target Information

**Gene name** ACTB

**Protein Name** Actin cytoplasmic 1

Organism	Gene ID	UniProt ID
Human	<a href="#">60;</a>	<a href="#">P60709;</a>

**Cellular Localization** Cytoplasm, cytoskeleton . Nucleus . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. .

**Tissue specificity** B-cell lymphoma,Brain,Cajal-Retzius cell,Eye,Fetal brain cortex,Foreskin,Hepatocellular car

**Function** Disease:Defects in ACTB are a cause of dystonia juvenile-onset (DYT) [MIM:607371]. DYT] is a form of dystonia with juvenile onset. Dystonia is defined by the presence of sustained involuntary muscle contraction, often leading to abnormal postures. DYT] patients manifest progressive, generalized, dopa-unresponsive dystonia, developmental malformations and sensory hearing loss.,Function:Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.,miscellaneous:In vertebrates 3 main groups of actin isoforms, alpha, beta and gamma have been identified. The alpha actins are found in muscle tissues and are a major constituent of the contractile apparatus. The beta and gamma actins coexist in most cell types as components of the cytoskeleton and as mediators of internal cell motility.,similarity:Belongs to the actin family.,subunit:Polymerization of globular actin (G-actin) leads to a structural filament (F-actin) in the form of a two-stranded helix. Each actin can bind to 4 others. Component of the BAF complex, which includes at least actin (ACTB), ARID1A, ARID1B/BAF250, SMARCA2, SMARCA4/BRG1, ACTL6A/BAF53, ACTL6B/BAF53B, SMARCE1/BAF57 SMARCC1/BAF155, SMARCC2/BAF170, SMARCB1/SNF5/INI1, and one or more of SMARCD1/BAF60A, SMARCD2/BAF60B, or SMARCD3/BAF60C. In muscle cells, the BAF complex also contains DPF3. Found in a complex with XPO6, Ran, ACTB and PFN1. Interacts with XPO6.,

## Validation Data

## Contact information

Orders: order.cn@immunoway.com  
Support: support.cn@immunoway.com  
Telephone: 408-747-0189 (USA) 400-8787-807(China)  
Website: <http://www.immunoway.com.cn>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**β-actin Monoclonal Antibody(5B7), Biotin Conjugated**

