

COX IV (6C8) Mouse mAb

CatalogNo: YM3033 **Orthogonal Validated** 

Key Features

Host Species

- Mouse

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF

MW

- 15kD (Observed)

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation PBS, pH 7.4, containing 0.5% BSA, 0.02% sodium azide as Preservative and 50% Glycerol.

Recommended Dilution Ratios

WB 1:1000-3000

IF 1:200

IHC 1:50-300

Basic Information

Clonality Monoclonal

Clone Number 6C8

Immunogen Information

Immunogen Recombinant Protein of Cytochrome c oxidase subunit 4 isoform 1, mitochondrial

Specificity The antibody detects endogenous COX IV protein.

Target Information

Gene name COX4I1

Protein Name Cytochrome c oxidase subunit 4 isoform 1, mitochondrial


Organism	Gene ID	UniProt ID
Human	1327;	P13073;
Mouse	12857;	P19783;
Rat	29445;	P10888;

Cellular Localization Mitochondrion inner membrane ; Single-pass membrane protein .


Tissue specificity Ubiquitous.

Function Function:This protein is one of the nuclear-coded polypeptide chains of cytochrome c oxidase , the terminal oxidase in mitochondrial electron transport. ,similarity:Belongs to the cytochrome c oxidase IV family. ,tissue specificity:Ubiquitous. ,


Validation Data




Zhang, Yanqiong, et al. "A discovery of clinically approved formula FBRP for repositioning to treat HCC by inhibiting PI3K/AKT/NF-κB activation." *Molecular Therapy-Nucleic Acids* 19 (2020): 890-904.



Western blot analysis of lysates from 1) COS7, 2) 3T3, 3) HeLa cells, (Green) primary antibody was diluted at 1:1000, 4°C over night, Dylight 800 secondary antibody (Immunoway:RS23910) was diluted at 1:10000, 37° 1hour. (Red) Actin β Polyclonal Antibody (Immunoway:YT0099) antibody was diluted at 1:5000 as loading control, 4°C over night, Dylight 680 secondary antibody (Immunoway:RS23720) was diluted at 1:10000, 37° 1hour.



Immunohistochemical analysis of paraffin-embedded Rat-kidney tissue.
 1,COX IV Monoclonal Antibody (6C8) was diluted at 1:200 (4°C overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C,20min). 3,Secondary antibody was diluted at 1:200 (room temperature, 30min). Negative control was used by secondary antibody only.



The nature compound dehydrocrenatidine exerts potent antihepatocellular carcinoma by destroying mitochondrial complexes in vitro and in vivo 2022 Feb 02. WB Human

Contact information

Orders: order.cn@immunoway.com
 Support: support.cn@immunoway.com
 Telephone: 400-8787-807(China)
 Website: <http://www.immunoway.com.cn>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
COX IV (6C8) Mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)