

#### www.immunoway.com.cn

# **CHSTB Rabbit pAb**

CatalogNo: YN0545

## Key Features

Host SpeciesRabbit

38kD (Observed)

MW

Reactivity
• Human,Mouse,Rat
Isotype

• IgG

ApplicationsWB,ELISA

#### **Recommended Dilution Ratios**

WB 1:500-2000 ELISA 1:5000-20000

### **Storage**

| Storage*    | -15°C to -25°C/1 year(Do not lower than -25°C)                          |
|-------------|---|
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |

#### **Basic Information**

Clonality Polyclonal

#### Immunogen Information

 Immunogen
 Synthesized peptide derived from part region of human protein


**Specificity** CHSTB Polyclonal Antibody detects endogenous levels of protein.

# Target Information

Gene name CHST11

| Protein Name             | Carbohydrate sulfotransferase 11 (Chondroitin 4-O-sulfotransferase 1) (Chondroitin 4-<br>sulfotransferase 1) (C4S-1) (C4ST-1) (C4ST1)  |               |                |  |
|--------------------------|--|---------------|----------------|--|
|                          | Organism   | Gene ID       | UniProt ID     |  |
|                          | Human  | <u>50515;</u> | <u>Q9NPF2;</u> |  |
|                          | Mouse  |               | <u>Q9JME2;</u> |  |
|                          | Rat  |               | <u>P69478;</u> |  |
| Cellular<br>Localization | Golgi apparatus membrane ; Single-pass type II membrane protein .  |               |                |  |
| Tissue specificity       | Widely expressed. Highly expressed in spleen, thymus, bone marrow, peripheral blood<br>leukocytes, lymph node, heart, brain, lung and placenta.  |               |                |  |
| Function                 | Catalytic activity:3'-phosphoadenylyl sulfate + chondroitin = adenosine 3',5'-bisphosphate<br>+ chondroitin 4'-sulfate.,Disease:A chromosomal aberration involving CHST11 is found in B-<br>cell chronic lymphocytic leukemias. Translocation t(12;14)(q23;q32) with<br>IgH.,Function:Catalyzes the transfer of sulfate to position 4 of the N-acetylgalactosamine<br>(GalNAc) residue of chondroitin. Chondroitin sulfate constitutes the predominant<br>proteoglycan present in cartilage and is distributed on the surfaces of many cells and<br>extracellular matrices. Can also sulfate Gal residues in desulfated dermatan sulfate.<br>Preferentially sulfates in GlcA->GalNAc unit than in IdoA->GalNAc unit. Does not form 4, 6-<br>di-O-sulfated GalNAc when chondroitin sulfate C is used as an acceptor.,PTM:N-<br>glycosylated; required for activity and stability.,similarity:Belongs to the sulfotransferase 2<br>family.,tissue specificity:Widely expressed. Highly expressed in spleen, thymus, bone<br>marrow, peripheral blood leukocytes, lymph node, heart, brain, lung and placenta., |               |                |  |

## **Validation Data**



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4°over night

# **Contact information**

Orders:order.cn@immunoway.comSupport:support.cn@immunoway.comTelephone:400-8787-807(China)Website:http://www.immunoway.com.cnAddress:2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: CHSTB Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents