

MUC4 Rabbit pAb

CatalogNo: YN0879

Key Features

Host Species

- Rabbit

Reactivity

- Human,Rat,Mouse,

Applications

- WB,ELISA

MW

- 238kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-2000

ELISA 1:5000-20000

Storage

Storage*

-15°C to -25°C/1 year(Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

Basic Information

Clonality

Polyclonal

Immunogen Information

Immunogen

Synthesized peptide derived from human protein . at AA range: 1970-2050

Specificity

MUC4 Polyclonal Antibody detects endogenous levels of protein.

Target Information

Gene name

MUC4

Protein Name	Mucin-4 (MUC-4) (Ascites sialoglycoprotein) (ASGP) (Pancreatic adenocarcinoma mucin) (Testis mucin) (Tracheobronchial mucin) [Cleaved into: Mucin-4 alpha chain (Ascites sialoglycoprotein 1) (ASGP-1); Mucin-4 beta chain (Ascites sialoglycoprotein 2) (ASGP-2)]		
	Organism	Gene ID	UniProt ID
	Human	4585;	Q99102;
	Mouse		Q8JZM8;
	Rat		Q63661;
Cellular Localization	[Mucin-4 beta chain]: Cell membrane ; Single-pass membrane protein . Isoforms lacking the Cys-rich region, EGF-like domains and transmembrane region are secreted. Secretion occurs by splicing or proteolytic processing. .; [Mucin-4 alpha chain]: Cell membrane . Secreted . Forms a complex with Mucin-4 beta chain at the cell membrane. .; [Isoform 3]: Cell membrane ; Single-pass membrane protein .; [Isoform 11]: Secreted .; [Isoform 15]: Secreted .; [Isoform 17]: Cell membrane ; Single-pass membrane protein .		
Tissue specificity	Expressed in the thymus, thyroid, lung, trachea, esophagus, stomach, small intestine, colon, testis, prostate, ovary, uterus, placenta, and mammary and salivary glands. Expressed in carcinomas arising from some of these epithelia, such as lung cancers, squamous cell carcinomas of the upper aerodigestive tract, mammary carcinomas, biliary tract, colon, and cervix cancers. Minimally or not expressed in the normal pancreas or chronic pancreatitis, but is highly expressed in pancreatic tumors and pancreatic tumor cell lines.		

Function

Alternative products: Additional isoforms exist, developmental stage: Expressed early in the primitive gut before respiratory and digestive epithelial cells have acquired their tissue and cell specificity. Expressed at the basal surface of the epithelium from week 14 to 26 weeks and then predominantly localized in only parietal cells. Immediately before birth, found in the cytoplasm of the mucous columnar epithelial cells. In the embryo expressed in skin, then disappears late in gestation., Function: May play a role in tumor progression. Ability to promote tumor growth may be mainly due to repression of apoptosis as opposed to proliferation. Has anti-adhesive properties. Seems to alter cellular behavior through both anti-adhesive effects on cell-cell and cell-extracellular matrix interactions and in its ability to act as an intramembrane ligand for ERBB2. Plays an important role in cell proliferation and differentiation of epithelial cells by inducing specific phosphorylation of ERBB2. The MUC4-ERBB2 complex causes site-specific phosphorylation of the ERBB2 'Tyr-1248'. In polarized epithelial cells segregates ERBB2 and other ERBB receptors and prevents ERBB2 from acting as a coreceptor. The interaction with ERBB2 leads to enhanced expression of CDKN1B. The formation of a MUC4-ERBB2-ERBB3-NRG1 complex leads to down-regulation of CDKN1B, resulting in repression of apoptosis and stimulation of proliferation., miscellaneous: Expression is a very useful predictor of poor prognosis in patients with invasive ductal carcinoma and intrahepatic cholangiocarcinoma, mass forming type (IDC, ICC-MF). Patients with IDC or ICC-MF who have high MUC4 expression had a worse survival rate than those with low MUC4 expression., PTM: Mucin-4 alpha chain is highly O-glycosylated., PTM: mucin-4 beta chain is predominantly N-glycosylated., PTM: Proteolytically cleaved into 2 chains, mucin-4 alpha chain and mucin-4 beta chain., sequence Caution: May be derived from an intron translation., similarity: Contains 1 AMOP domain., similarity: Contains 1 NIDO domain., similarity: Contains 1 VWFD domain., similarity: Contains 2 EGF-like domains., subcellular location: Isoforms lacking the Cys-rich region, EGF-like domains and transmembrane region are secreted. Secretion occurs by splicing or proteolytic processing., subunit: A heterodimeric complex, composed of a mucin-4 alpha chain and a cysteine-rich transmembrane mucin-4 beta chain. Mucin-4 beta chain interacts with ERBB2 via the EGF-like domain 1. In nonpolarized cells, associates with ERBB2 and ERBB3., tissue specificity: Expressed in the thymus, thyroid, lung, trachea, esophagus, stomach, small intestine, colon, testis, prostate, ovary, uterus, placenta, and mammary and salivary glands. Expressed in carcinomas arising from some of these epithelia, such as lung cancers, squamous cell carcinomas of the upper aerodigestive tract, mammary carcinomas, biliary tract, colon, and cervix cancers. Minimally or not expressed in the normal pancreas or chronic pancreatitis, but is highly expressed in pancreatic tumors and pancreatic tumor cell lines.,

| Validation Data

| Contact information

Orders: order.cn@immunoway.com
Support: support.cn@immunoway.com
Telephone: 400-8787-807(China)
Website: <http://www.immunoway.com.cn>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
MUC4 Rabbit pAb

