

DRA Rabbit pAb

CatalogNo: YN2348

Key Features

Host Species

Rabbit

Reactivity
• Human

ApplicationsWB,ELISA

Maddi

MW

Isotype

27kD (Observed)

• IgG

Recommended Dilution Ratios

WB 1:500-2000

ELISA 1:5000-20000

Storage

Storage*

-15°C to -25°C/1 year(Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

Basic Information

Clonality

Polyclonal

Immunogen Information

Immunogen

Synthesized peptide derived from human protein . at AA range: 100-180

Specificity

DRA Polyclonal Antibody detects endogenous levels of protein.

| Target Information

Gene name

HLA-DRA HLA-DRA1

Protein Name

HLA class II histocompatibility antigen, DR alpha chain (MHC class II antigen DRA)

| Organism | Gene ID | UniProt ID | |
|----------|---------------|-----------------|--|
| Human | <u>3122</u> ; | <u>P01903</u> ; | |

Cellular Localization

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein. Early endosome membrane: Single-pass type I membrane protein. Late endosome membrane; Single-pass type I membrane protein. Lysosome membrane; Single-pass type I membrane protein. Autolysosome membrane; Single-pass type I membrane protein. The MHCII complex transits through a number of intracellular compartments in the endocytic pathway until it reaches the cell membrane for antigen presentation (PubMed:9075930, PubMed:18305173). Component of immunological synapses at the interface between T cell and APC (PubMed:15322540, PubMed:29884618). .

Tissue specificity Expressed in professional APCs: macrophages, dendritic cells and B cells (at protein level) (PubMed:31495665, PubMed:15322540, PubMed:23783831). Expressed in thymic epithelial cells (at protein level) (PubMed:23783831).

Function

Disease:Genetic variations in HLA-DRA are associated with susceptibility to hepatitis B virus infection (HBV infection) [MIM:610424]. Approximately one third of all cases of cirrhosis and half of all cases of hepatocellular carcinoma can be attributed to chronic HBV infection. HBV infection may result in subclinical or asymptomatic infection, acute self-limited hepatitis, or fulminant hepatitis requiring liver transplantation.,polymorphism:The following alleles of DRA are known: DRA*0101 and DRA*0102. The sequence shown is that of

DRA*0101., similarity: Belongs to the MHC class II family., similarity: Contains 1 Ig-like C1-type (immunoglobulin-like) domain., subunit: Heterodimer of an alpha chain and a beta chain.,

Validation Data

I Contact information

Orders: order.cn@immunoway.com Support: support.cn@immunoway.com

Telephone: 400-8787-807(China)

Website: http://www.immunoway.com.cn

2200 Ringwood Ave San Jose, CA 95131 USA Address:



Please scan the OR code to access additional product information: DRA Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents