

ADAMTS-16 Rabbit pAb

CatalogNo: YT0114

Key Features

| Host Species • Rabbit | Reactivity Human,Mouse | Applications IHC,IF,ELISA |
|----------------------------|--|---|
| MW • 136kD (Calculated) | Isotype • IgG | |

Recommended Dilution Ratios

IHC 1:100-1:300 ELISA 1:40000 IF 1:50-200

Storage

Storage*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

ImmunogenSynthesized peptide derived from ADAMTS-16 . at AA range: 440-520SpecificityADAMTS-16 Polyclonal Antibody detects endogenous levels of ADAMTS-16 protein.

Target Information

Gene name ADAMTS16

Protein Name

A disintegrin and metalloproteinase with thrombospondin motifs 16

| Organism | Gene ID | UniProt ID |
|----------|----------------|----------------|
| Human | <u>170690;</u> | <u>Q8TE57;</u> |
| Mouse | <u>271127;</u> | <u>Q69Z28;</u> |


Cellular Secreted, extracellular space, extracellular matrix.

Localization

Tissue specificity Expressed in fetal lung and kidney and in adult prostate and ovary.

Function cofactor:Binds 1 zinc ion per subunit.,Domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme., Domain: The spacer domain and the TSP type-1 domains are important for a tight interaction with the extracellular matrix., PTM: The precursor is cleaved by a furin endopeptidase., similarity: Contains 1 disintegrin domain., similarity: Contains 1 peptidase M12B domain., similarity: Contains 1 PLAC domain., similarity: Contains 6 TSP type-1 domains.,tissue specificity:Expressed in fetal lung and kidney and in adult prostate and ovary.,

Validation Data



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 30min).

Contact information

| Orders: | order.cn@immunoway.com |
|------------|--|
| Support: | support.cn@immunoway.com |
| Telephone: | 400-8787-807(China) |
| Website: | http://www.immunoway.com.cn |
| Address: | 2200 Ringwood Ave San Jose, CA 95131 USA |



Please scan the QR code to access additional product information: ADAMTS-16 Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents