

www.immunoway.com.cn

NGF Rabbit pAb

CatalogNo: YT3114 Orthogonal Validated 💽

Key Features

Host Species Rabbit 	Reactivity Human,Mouse,Rat 	Applications • WB,IHC,IF,ELISA
MW • 27kD (Observed)	Isotype • IgG	

Recommended Dilution Ratios

WB 1:500-1:2000 IHC: 1:100-300 ELISA 1:20000 IF 1:100-300 Not yet tested in other applications.

Storage

Storage*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

ImmunogenThe antiserum was produced against synthesized peptide derived from human NGF. AA
range:33-82SpecificityNGF Polyclonal Antibody detects endogenous levels of NGF protein.

Target Information

Gene	name	NGF
------	------	-----

Protein Name Beta-nerve growth factor


Organism	Gene ID	UniProt ID
Human	<u>4803;</u>	<u>P01138;</u>
Mouse	<u>18049;</u>	<u>P01139;</u>
Rat	<u>310738;</u>	<u>P25427;</u>

CellularSecreted . Endosome lumen . ProNGF is endocytosed after binding to the cell surfaceLocalizationreceptor formed by SORT1 and NGFR. .


Tissue specificity Brain, Epithelium, Eye, Leukocyte,


FunctionDisease:Defects in NGF are the cause of hereditary sensory and autonomic neuropathy type
5 (HSAN5) [MIM:608654]. The hereditary sensory and autonomic neuropathies are a
genetically and clinically heterogeneous group of disorders characterized by degeneration
of dorsal root and autonomic ganglion cells, and by sensory and/or autonomic
abnormalities. HSAN5 patients manifest loss of pain perception and impaired temperature
sensitivity, ulcers, and in some cases self-mutilation. The autonomic involvement is
variable.,Function:Nerve growth factor is important for the development and maintenance
of the sympathetic and sensory nervous systems. It stimulates division and differentiation
of sympathetic and embryonic sensory neurons.,online information:Nerve growth factor
entry,similarity:Belongs to the NGF-beta family.,subunit:Homodimer.,

Validation Data



Immunofluorescence analysis of A549. 1,primary Antibody(red) was diluted at 1:200(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 594 Secondary antibody was diluted at 1:1000(room temperature, 50min).3, DAPI(blue) 10min.





Western Blot analysis of Jurkat cells using NGF Polyclonal Antibody diluted at $1{:}2000$

Immunohistochemistry analysis of NGF antibody in paraffin-embedded human lung carcinoma tissue.

Western blot analysis of lysate from Jurkat cells, using NGF antibody.

Contact information

Orders:	order.cn@immunoway.com
Support:	support.cn@immunoway.com
Telephone:	400-8787-807(China)
Website:	http://www.immunoway.com.cn
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **NGF Rabbit pAb** For Research Use Only. Not for Use in Diagnostic Procedures.

Immunoway - 4 / 4