

www.immunoway.com.cn

IP3R-I Rabbit pAb

CatalogNo: YT5663

Key Features

Host Species Rabbit 	Reactivity Human,Mouse,Rat 	App • W
MW • 320kD (Observed)	Isotype • IgG	

Applications
• WB,IHC

Recommended Dilution Ratios

WB 1:500-2000 IHC 1:50-300

Storage

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen	Synthesized peptide derived from IP3R-I . at AA range: 1700-1780
Specificity	IP3R-I Polyclonal Antibody detects endogenous levels of IP3R-I protein.

Target Information

Gene name ITPR1

Protein Name Inositol 1,4,5-trisphosphate receptor type 1

Organism	Gene ID	UniProt ID
Human	<u>3708;</u>	<u>Q14643;</u>
Mouse	<u>16438;</u>	<u>P11881;</u>
Rat	<u>25262;</u>	<u>P29994;</u>


Cellular Localization

Endoplasmic reticulum membrane ; Multi-pass membrane protein . Cytoplasmic vesicle, secretory vesicle membrane ; Multi-pass membrane protein . Cytoplasm, perinuclear region . Endoplasmic reticulum and secretory granules (By similarity). .


Tissue specificity Widely expressed.

Function Alternative products: There is a combination of three alternatively spliced domains at site SI, SIII and site SII (A and C). Experimental confirmation may be lacking for some isoforms, Disease: Defects in ITPR1 are the cause of spinocerebellar ataxia type 15 (SCA15) (SCA15) [MIM:606658]. Spinocerebellar ataxia is a clinically and genetically heterogeneous group of cerebellar disorders. Patients show progressive incoordination of gait and often poor coordination of hands, speech and eye movements, due to degeneration of the cerebellum with variable involvement of the brainstem and spinal cord. SCA15 is an autosomal dominant cerebellar ataxia (ADCA). It is very slow progressing form with a wide range of onset, ranging from childhood to adult. Most patients remain ambulatory..Domain:The receptor contains a calcium channel in its C-terminal extremity. Its large N-terminal cytoplasmic region has the ligand-binding site in the N-terminus and modulatory sites in the middle portion immediately upstream of the channel region.,Function:Intracellular channel that mediates calcium release from the endoplasmic reticulum following stimulation by inositol 1.4.5-trisphosphate..miscellaneous:Calcium appears to inhibit ligand binding to the receptor, most probably by interacting with a distinct calcium-binding protein which then inhibits the receptor., PTM: Phosphorylated by cAMP kinase. Phosphorylation prevents the ligand-induced opening of the calcium channels., PTM: Phosphorylated on tyrosine residues., similarity: Belongs to the InsP3 receptor family., similarity: Contains 5 MIR domains., subunit: Homotetramer. Interacts with TRPC4. The PPXXF motif binds HOM1. HOM2 and HOM3. Interacts with RYR1. RYR2. ITPR1. SHANK1 and SHANK3. Interacts with ERP44 in a pH-, redox state- and calcium-dependent manner which results in the inhibition the calcium channel activity. The strength of this interaction inversely correlates with calcium concentration. Part of cGMP kinase signaling complex at least composed of ACTA2/alpha-actin, CNN1/calponin H1, PLN/phospholamban, PRKG1 and ITPR1. Interacts with AHCYL1 (By similarity). Interacts with MRVI1., tissue specificity: Widely expressed.,

Validation Data



Western blot analysis of 293T using IP3R-I antibody. Antibody was diluted at 1:500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Contact information

Orders:	order.cn@immunoway.com
Support:	support.cn@immunoway.com
Telephone:	400-8787-807(China)
Website:	http://www.immunoway.com.cn
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **IP3R-I Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents