

#### www.immunoway.com.cn

# CD86 Rabbit pAb

CatalogNo: YT7823 Orthogonal Validated 💽

# Key Features

| Host Species <ul> <li>Rabbit</li> </ul> | Reactivity<br>• Human,Mouse | <ul><li>Applications</li><li>WB,IF,ELISA</li></ul> |
|---|-----------------------------|--|
| MW<br>• 60-80kD (Calculated)            | Isotype<br>• IgG            |  |

### **Recommended Dilution Ratios**

WB 1:500-2000 IF/ICC1:100-500 ELISA 1:5000-20000

### **Storage**

| Storage*    | -15°C to -25°C/1 year(Do not lower than -25°C)                          |
|-------------|---|
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |

### **Basic Information**

Clonality Polyclonal

### Immunogen Information

| Immunogen   | Synthesized peptide derived from human CD86           |
|-------------|---|
| Specificity | This antibody detects endogenous levels of Human CD86 |

# Target Information

Gene name CD86 CD28LG2

#### Protein Name CD86

| Organism | Gene ID       | UniProt ID     |
|----------|---------------|----------------|
| Human    | <u>942;</u>   | <u>P42081;</u> |
| Mouse    | <u>12524;</u> | <u>P42082;</u> |

Cellular Cell membrane; Single-pass type I membrane protein.

Localization

**Tissue specificity** Expressed by activated B-lymphocytes and monocytes.

Function Function:Receptor involved in the costimulatory signal essential for T-lymphocyte proliferation and interleukin-2 production, by binding CD28 or CTLA-4. May play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation. Isoform 2 interferes with the formation of CD86 clusters, and thus acts as a negative regulator of T-cell activation.,online information:CD86 entry,PTM:Polyubiquitinated; which is promoted by MARCH8 and results in endocytosis and lysosomal degradation.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,subunit:Interacts with MARCH8. Interacts with human herpesvirus 8 MIR2 protein (Probable). Interacts with adenovirus subgroup B fiber proteins and acts as a receptor for these viruses.,tissue specificity:Expressed by activated B-lymphocytes and monocytes.,

# Validation Data

## **Contact information**

| Orders:    | order.cn@immunoway.com                   |
|------------|--|
| Support:   | support.cn@immunoway.com                 |
| Telephone: | 400-8787-807(China)                      |
| Website:   | http://www.immunoway.com.cn              |
| Address:   | 2200 Ringwood Ave San Jose, CA 95131 USA |
|            |  |



Please scan the QR code to access additional product information: **CD86 Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents